Molecular Ecology (2002) **11**, 2717–2728

Author index

Abatzopoulos TJ see Triantafyllidis A
Abbot P, and Moran NA. Extremely low levels of genetic polymorphism in endosymbionts (*Buchnera*) of aphids (*Pemphigus*), 2649
Abbott RJ see Holderegger R
Abebe E see Floyd R
Achituv Y see Appelbaum L
Adamowicz SJ, Ryan Gregory T, Marinone MC & Hebert PDN. New insights into the distribution of polyploid *Daphnia*: the Holarctic revisited and Argentina explored, 1209
Adams RJ see Johnson KP
Adams RMM see Green AM
Adams WT see Burczyk J
Afzal-Rafii Z see Dodd RS
Aide TM see Rivera-Ocasio E
Åkesson S see Bensch S
Albon SD see Côté SD
Allendorf FW see Tallmon DA
Alpers D see Joseph L
Amarista M see Mavárez J
Ambrosini R see Saino N
Amos W see Johnson PCD
Amos W see Valsecchi E
Andreone F see Vences M
Angers B see Charbonnel N
Angulo E & Cooke B. First synthesize new viruses then regulate their release? The case of the wild rabbit, 2703
Appelbaum L, Achituv Y & Mokady O. Speciation and the establishment of zonation in an intertidal barnacle: specific settlement vs. selection, 1731
Apsit VJ see Sork VL
Apte S & Gardner JPA. Population genetic subdivision in the New Zealand greenshell mussel (*Perna canaliculus*) inferred from single-strand conformation polymorphism analysis of mitochondrial DNA, 1617
Arafeh RMH, Sapir Y, Shmida A, Iraki N, Fragman O & Comes HP. Patterns of genetic and phenotypic variation in *Iris haynei* and *I. atrofusca* (*Iris* sect. *Oncocyclus* = the royal irises) along an eco-geographical gradient in Israel and the West Bank, 39
Arctander P see Pitra C
Aron S see Fournier D
Arroyo García R, Martínez Zapater JM, García Criado B & Zabalgogeazcoa I. Genetic structure of natural populations of the grass endophyte *Epichloë festucae* in semiarid grasslands, 355
Asoh K see McCafferty S
Atagi DY see Heath DD
Atkinson HJ see Cowgill SE
Atkinson RJ see Rokas A
Austerlitz F see Mariette S
Avise JC see Fiumera AC
Avise JC see MacKiewicz M
Avise JC see Walker D
Ayasse M see Paxton RJ
Ayre DJ see England PR
Bachmann K see Stehlik I
Bacles C see Bonnin I
Ball Jr RM see Moore MK
Ballment E see Benzie JAH
Balloux F see Fumagalli L
Balloux F & Goudet J. Statistical properties of population differentiation estimators under stepwise mutation in a finite island model, 771
Balloux F & Lugon-Moulin N. The estimation of population differentiation with microsatellite markers, 155
Barber PH, Palumbi SR, Erdmann MV & Moosa MK. Sharp genetic breaks among populations of *Haptosquilla pulchella* (Stomatopoda) indicate limits to larval transport: patterns, causes, and consequences, 659
Barfuss M see Schönswetter P
Bargelloni L see Zane L
Baric S see Salzburger W
Barlow M see Dawson MN
Barrett JA see Trewick SA
Barth S, Melchinger AE & Lübberstedt Th. Genetic diversity in *Arabidopsis thaliana* L. Heynh. investigated by cleaved amplified polymorphic sequence (CAPS) and inter-simple sequence repeat (ISSR) markers, 495
Bass AL see Rocha LA
Bassano B see Maudet C
Bateman RM see Squirrell J
Battisti A see Salvato P
Beauwens T see Vekemans X
Beiles A see Li Y-C
Bekkevold D see Hansen MM
Bekkevold D, Hansen MM & Loeschcke V. Male reproductive competition in spawning aggregations of cod (*Gadus morhua*, L.), 91
Ben-David M see Blundell GM
Bensch S see Waldenström J
Bensch S, Åkesson S & Irwin DE. The use of AFLP to find an informative SNP: genetic differences across a migratory divide in willow warblers, 2359
Bensch S, Helbig AJ, Salomon M & Seibold I. Amplified fragment length polymorphism analysis identifies hybrids between two subspecies of warblers, 473
Bentley JK see Jiggins FM
Bentzen P see Kinnison MT
Bentzen P see Larson S
Benzie JAH, Ballment E, Forbes AT, Demetriades NT, Sugama K, Haryanti & Moria S. Mitochondrial DNA variation in Indo-Pacific populations of the giant tiger prawn, *Penaeus monodon*, 2553
Bermingham E see McCafferty S
Bernatchez L see Dufresne F
Bernini F see Salomone N
Berrebi P see Fumagalli L
Bertorelle G see Vernesi C
Betancourt JL see Kuch M
Bidartondo MI & Bruns TD. Fine-level mycorrhizal specificity in the Monotropoideae (Ericaceae): specificity for fungal species groups, 557

© 2002 Blackwell Science Ltd

Biémont C see Lepetit D
Blackburn C see Dallimer M
Blattner FR see Stehlik I
Blaxter M see Floyd R
Blundell GM, Ben-David M, Groves P, Bowyer RT & Geffen E. Characteristics of sex-biased dispersal and gene flow in coastal river otters: implications for natural recolonization of extirpated populations, 289
Boitani L see Lucchini V
Bonhomme F see Meusnier I
Bonizzoni M, Katsoyannos BI, Marguerie R, Guglielmino CR, Gasperi G, Malacrida A & Chapman T. Microsatellite analysis reveals remating by wild Mediterranean fruit fly females, *Ceratitis capitata*, 1915
Bonnin I, Colas B, Bacles C, Holl A-C, Hendoux F, Destiné B & Viard F. Population structure of an endangered species living in contrasted habitats: *Parnassia palustris* (Saxifragaceae), 979
Borge T see Primmer CR
Bougrier S see Garnier-Géré PH
Bourget E see Dufresne F
Bowen BW see Rocha LA
Bower RT see Blundell GM
Brachet S see Morand M-E
Brachet S see Morand-Prieur ME
Brehm A see Lepetit D
Breitenmoser-Würsten C see Maudet C
Bremond P see Charbonnel N
Brochmann C see Gabrielsen TM
Brodie III ED see Janzen FJ
Brodie Jr ED see Janzen FJ
Brookfield JFY see MacDonald C
Brossier F see Fumagalli L
Brown AHD see Rauscher JT
Brown JK see Legg JP
Brown M see Waldick RC
Brown S see Colgan D
Bruford MW see Paulo OS
Bruns TD see Bidartondo MI
Brunsfeld SJ see Richardson BA
Brzosko E, Wróblewska A & Ratkiewicz M. Spatial genetic structure and clonal diversity of island populations of lady's slipper (*Cypripedium calceolus*) from the Biebrza National Park (northeast Poland), 2499
Bucci G, & Menozzi P. Spatial autocorrelation and linkage of Mendelian RAPD markers in a population of *Picea abies* Karst., 305
Budrick J see Dodd RS
Bull CM see Gardner MG
Burczyk J, Adams WT, Moran GF & Griffin AR. Complex patterns of mating revealed in a *Eucalyptus regnans* seed orchard using allozyme markers and the neighbourhood model, 2379
Burden JP, Griffiths CM, Cory JS, Smith P & Sait SM. Vertical transmission of sublethal granulovirus infection in the Indian meal moth, *Plodia interpunctella*, 547
Burke T see Châline N
Burton C, Krebs CJ & Taylor EB. Population genetic structure of the cyclic snowshoe hare (*Lepus americanus*) in southwestern Yukon, Canada, 1689
Busch C see Heath DD
Bush GL see Weiblen GD
Cabrera H see Doums C
Caccone A see Ciofi C
Calza S see Saino N
Caramelli D see Vernesi C
Cariello MO, Schwabb HG, Lee RW & Macedo RHF. Individual female clutch identification through yolk protein electrophoresis in the communally-breeding guira cuckoo (*Guira guira*), 2417
Carvalho GR see Hänfling B
Casais CA see Moyes CL
Cassis G see Colgan D
Castilla JC, Collins AG, Meyer CP, Guiñez R & Lindberg DR. Recent introduction of the dominant tunicate, *Pyura praeputialis* (Urochordata, Pyuridae) to Antofagasta, Chile, 1579
Châline N, Ratnieks FLW & Burke T. Anarchy in the UK: detailed genetic analysis of worker reproduction in a naturally occurring British anarchistic honeybee, *Apis mellifera*, colony using DNA microsatellites, 1795
Chambon J-M see Kerdelhué C
Chapman T see Bonizzoni M
Charbonnel N, Angers B, Rasatavonjizay R, Bremond P, Debain C & Jarne P. The influence of mating system, demography, parasites and colonization on the population structure of *Biomphalaria pfeifferi* in Madagascar, 2213
Charlwood JD see Pinto J
Chevre AM see Gueritaine G
Chorlton KH see Sackville Hamilton NR
Chorlton KH see Skøt L
Christie F see Colgan D
Churikov D & Gharrett AJ. Comparative phylogeography of the two pink salmon broodlines: an analysis based on a mitochondrial DNA genealogy, 1077
Cicero C see Johnson NK
Ciofi C, Milinkovitch MC, Gibbs JP, Caccone A & Powell JR. Microsatellite analysis of genetic divergence among populations of giant Galápagos tortoises, 2265
Clauss MJ, Cobban H & Mitchell-Olds T. Cross-species microsatellite markers for elucidating population genetic structure in *Arabidopsis* (Brassicacae), 591
Clayton DH see Johnson KP
Clegg SM see Kimura M
Cobban H see Clauss MJ
Cogan R see Williams CL
Colas B see Bonnin I
Cole SG see Moyes CL
Colgan D, Brown S, Major RE, Christie F, Gray MR & Cassis G. Population genetics of wolf spiders of fragmented habitat in the wheat belt of New South Wales, 2295
Collins AG see Castilla JC
Colson I. Selection and gene flow between microenvironments: the case of *Drosophila* at Lower Nahal Oren, Mount Carmel, Israel, 1311
Comes HP see Arafeh RMH
Comes HP, Kropf M & Kadereit JW. Late Quaternary distributional stasis in the submediterranean mountain plant *Anthyllis montana* L. (Fabaceae) inferred from ITS sequences and AFLP markers, 447
Comstock KE, Georgiadis N, Pecon-Slattery J, Roca AL Ostrander EA, O'Brien SJ & Wasser SK. Patterns of molecular genetic variation among African elephant populations, 2489
Concato S see Salvato P

Consuegra S, García De Léaniz C, Serdio A, González Morales M, Straus LG, Knox D & Verspoor E. Mitochondrial DNA variation in Pleistocene and modern Atlantic salmon from the Iberian glacial refugium, 2037
Cook JA see Fleming MA
Cook JA see Stone KD
Cooke B see Angulo E
Cooper SJB see Gardner MG
Cooper SJB see Ibrahim KM
Corkeron P see Valsecchi E
Cornuet J-M see Estoup A
Cory JS see Burden JP
Côté SD, Dallas JF, Marshall F, Irvine RJ, Langvatn R & Albon SD. Microsatellite DNA evidence for genetic drift and philopatry in Svalbard reindeer, 1923
Cottin C see Triantafyllidis A
Coutinho TA see Slippers B
Cowgill SE, Wright C & Atkinson HJ. Transgenic potatoes with enhanced levels of nematode resistance do not have altered susceptibility to nontarget aphids, 821
Crespi BJ see Law JH
Crim JL, Spotila LD, Spotila JR, O'Connor M, Reina R, Williams CJ & Paladino FV. The leatherback turtle, *Dermochelys coriacea*, exhibits both polyandry and polygyny, 2097
Crivelli AJ see Fumagalli L
Croft JM see King RA

Dahlgaard J see Frydenberg J
Dale PJ see Moyes CL
Dallas JF see Côté SD
Dallimer M, Blackburn C, Jones PJ & Pemberton JM. Genetic evidence for male biased dispersal in the red-billed quelea *Quelea quelea*, 529
Daniell TJ see Vandenkoornhuyse P
Darmency H see Gueritaine G
Dávila JA see Potti J
Davis FW see Sork VL
Dawson MN, Louie KD, Barlow M, Jacobs DK & Swift CC. Comparative phylogeography of sympatric sister species, *Clevelandia ios* and *Eucyclogobius newberryi* (Teleostei, Gobiidae), across the California Transition Zone, 1065
De Meeûs T see Prugnolle F
Debain C see Charbonnel N
Degnan BM see Wörheide G
Delmotte F, Leterme N, Gauthier J-P, Rispe C & Simon J-C. Genetic architecture of sexual and asexual populations of the aphid *Rhopalosiphum padi* based on allozyme and microsatellite markers, 711
Demetriades NT see Benzie JAH
Desmarais E see Meusnier I
Despres L, Loriot S & Gaudeul M. Geographic pattern of genetic variation in the European globeflower *Trollius europaeus* L. (Ranunculaceae) inferred from amplified fragment length polymorphism markers, 2337
Destiné B see Bonnin I
Destombe C see Meusnier I
DeWoody JA see Mackiewicz M
Dey DK see Holsinger KE
Do Rosário VE see Pinto J
Dodd RS, Afzal-Rafii Z, Kashani N & Budrick J. Land barriers and open oceans: effects on gene diversity and population structure in *Avicennia germinans* L. (Avicenniaceae), 1327
Donnelly MJ see Pinto J
Douhan GW, Peever TL & Murray TD. Multilocus population structure of *Tapesia yallundae* in Washington State, 2229
Doums C, Cabrera H & Peeters C. Population genetic structure and male-biased dispersal in the queenless ant *Diacamma cyaneiventre*, 2251
Downie DA. Locating the sources of an invasive pest, grape phylloxera, using a mitochondrial DNA gene genealogy, 2013
Doyle JJ see Rauscher JT
Draheim HM see Tallmon DA
Drown DM see Johnson KP
Duck JM see Vandenkoornhuyse P
Dufour J see Morand M-E
Dufresne F, Bourget E & Bernatchez L. Differential patterns of spatial divergence in microsatellite and allozyme alleles: further evidence for locus-specific selection in the acorn barnacle, *Semibalanus balanoides*?, 113
Dupanloup I, Schneider S & Excoffier L. A simulated annealing approach to define the genetic structure of populations, 2571
Durand P see Prugnolle F
Duron O see Fumagalli L
Dutech C, Seiter J, Petronelli P, Joly HI & Jarne P. Evidence of low gene flow in a neotropical clustered tree species in two rainforest stands of French Guiana, 725
Dyer RJ see Sork VL

Eber F see Gueritaine G
Ehinger M, Fontanillas P, Petit E & Perrin N. Mitochondrial DNA variation along an altitudinal gradient in the greater white-toothed shrew, *Crocidura russula*, 939
Elissa N see Pinto J
Elliott NG see Jarman SN
Emerson BC see Salomone N
Emerson BC. Evolution on oceanic islands: molecular phylogenetic approaches to understanding pattern and process, 951
England PR, Usher AV, Whelan RJ & Ayre DJ. Microsatellite diversity and genetic structure of fragmented populations of the rare, fire-dependent shrub *Grevillea macleayana*, 967
Englbrecht CC, Schliewen U & Tautz D. The impact of stocking on the genetic integrity of Arctic charr (*Salvelinus*) populations from the Alpine region, 1017
Ennos RA see Rendell S
Eppe R see Weiss S
Erdmann MV see Barber PH
Estoup A, Jarne P & Cornuet J-M. Homoplasy and mutation model at microsatellite loci and their consequences for population genetics analysis, 1591
Etnier M see Larson S
Evans TA & Goodisman MAD. Nestmate relatedness and population genetic structure of the Australian social crab spider *Diaea ergandros* (Araneae: Thomisidae), 2307
Excoffier L see Dupanloup I

Fabbri E see Lucchini V
Fahima T see Li Y-C
Failloux A-B see Huber K
Felder DL see Mathews LM
Fernandez-M JF see Sork VL
Ferreira C see Pinto J

Field J see Paxton RJ

Finston T. Geographic patterns of population genetic structure in *Mytilocypris* (Ostracoda: Cyprididae): interpreting breeding systems, gene flow and history in species with differing distributions, 1931

Fitter AH see Vandenkoornhuyse P

Fiumera AC, Porter BA, Grossman GD & Avise JC. Intensive genetic assessment of the mating system and reproductive sucess in a semi-closed population of the mottled sculpin, *Cottus bairdi*, 2367

Fleming M see Larson S

Fleming MA & Cook JA. Phylogeography of endemic ermine (*Mustela erminea*) in southeast Alaska, 795

Flemming AF see Matthee CA

Fletcher DE see Mackiewicz M

Floyd R, Abebe E, Papert A & Blaxter M. Molecular barcodes for soil nematode identification, 839

Flynn RW see Stone KD

Fontanillas P see Ehinger M

Forbes AT see Benzie JAH

Ford BA see Jerome CA

Ford MJ. Applications of selective neutrality tests to molecular ecology, 1245

Forichon J see Kerdelhué C

Foster WA see Johnson PCD

Fouillet P see Lepetit D

Fournier D, Aron S & Milinkovitch MC. Investigation of the population genetic structure and mating system in the ant *Pheidole pallidula*, 1805

Fragman O see Arafeh RMH

Frascaria-Lacoste N see Morand M-E

Frascaria-Lacoste N see Morand-Prieur ME

Freeman-Gallant CR, Johnson EM, Saponara F & Stanger M. Variation at the major histocompatibility complex in Savannah sparrows, 1125

French R see Legg JP

Frías Ó see Potti J

Frydenberg J, Pertoldi C, Dahlgaard J & Loeschcke V. Genetic variation in original and colonizing *Drosophila buzzatii* populations analysed by microsatellite loci isolated with a new PCR screening method, 181

Fuertes Aguilar J see Gutiérrez Larena B

Fukatsu T see Kondo N

Fukatsu T see Tsuchida T

Fumagalli L, Snoj A, Jesenšek D, Balloux F, Jug T, Duron O, Brossier F, Crivelli AJ & Berrebi P. Extreme genetic differentiation among the remnant populations of marble trout (*Salmo marmoratus*) in Slovenia, 2711

Gabrielsen TM, Brochmann C & Rueness J. The Baltic Sea as a model system for studying postglacial colonization and ecological differentiation, exemplified by the red alga *Ceramium tenuicorne*, 2083

Gales NJ see Jarman SN

Gallery R see Husband R

Gan YY see Teo LL

Gantenbein B & Largiadèr CR. *Mesobuthus gibbosus* (Scorpiones: Buthidae) on the island of Rhodes — hybridization between Ulysses' stowaways and native scorpions?, 925

García Criado B see Arroyo García R

García De Léaniz C see Consuegra S

Gardner JPA see Apte S

Gardner MG, Bull CM & Cooper SJB. High levels of genetic monogamy in the group-living Australian lizard *Egernia stokesii*, 1787

Garnier-Géré PH, Naciri-Graven Y, Bougrier S, Magoulas A, Héral M, Kotoulas G, Hawkins A & Gérard A. Influences of triploidy, parentage and genetic diversity on growth of the Pacific oyster *Crassostrea gigas* reared in contrasting natural environments, 1499

Garrigan D see Hedrick PW

Gasparich GE see Meixner MD

Gasperi G see Bonizzoni M

Gaudeul M see Despres L

Gauthier D see Maudet C

Gauthier J-P see Delmotte F

Geffen E see Blundell GM

Georgiadis N see Comstock KE

Gérard A see Garnier-Géré PH

Gharrett AJ see Churikov D

Gibbs JP see Ciofi C

Gibby M see Trewick SA

Gibson G. Microarrays in ecology and evolution: a preview, 17

Gil V see Pinto J

Gill PC see Jarman SN

Girman DJ see Kimura M

Gjøstein H see Røed KH

Glaw F see Vences M

Godé C see Meusnier I

Godoy JA see Palomares F

González Morales M see Consuegra S

Goodacre SL. Population structure, history and gene flow in a group of closely related land snails: genetic variation in *Partula* from the Society Islands of the Pacific, 55

Goodisman MAD see Evans TA

Gornall RJ see King RA

Goudet J see Balloux F

Goudet J, Perrin N & Waser P. Tests for sex-biased dispersal using bi-parentally inherited genetic markers, 1103

Grady JM see Hardy ME

Gray MR see Colgan D

Green AM, Mueller UG & Adams RMM. Extensive exchange of fungal cultivars between sympatric species of fungus-growing ants, 191

Greenberg DL see Mock KE

Griffin AR see Burczyk J

Griffith SC, Owens IPF & Thuman, KA. Extra pair paternity in birds: a review of interspecific variation and adaptive function, 2195

Griffiths CM see Burden JP

Grivet D & Petit RJ. Phylogeography of the common ivy (*Hedera* sp.) in Europe: genetic differentiation through space and time, 1351

Groen AF see van Hooft WF

Gross R see Grunwald C

Grossman GD see Fiumera AC

Groves P see Blundell GM

Grunwald C, Stabile J, Waldman JR, Gross R & Wirgin I. Population genetics of shortnose sturgeon *Acipenser brevirostrum* based on mitochondrial DNA control region sequences, 1885

Gueritaine G, Sester M, Eber F, Chevre AM & Darmency H. Fitness of blackcross six of hybrids between transgenic oilseed rape (*Brassica napus*) and wild radish (*Raphanus raphanistrum*), 1419

Guglielmino CR see Bonizzoni M

Guiñez R see Castilla JC

Gutiérrez Larena B, Fuertes Aguilar J & Nieto Feliner G. Glacial-induced altitudinal migrations in *Armeria* (Plumbaginaceae) inferred from patterns of chloroplast DNA haplotype sharing, 1965

Guyomard R see Triantafyllidis A

Hadfield MG see Holland BS

Haeger PD see Moyes CL

Haldimann P see Steinger T

Hale P see Valsecchi E

Hänfling B, Hellemans B, Volckaert FAM & Carvalho GR. Late glacial history of the cold-adapted freshwater fish *Cottus gobio*, revealed by microsatellites, 1717

Hangelbroek HH, Ouborg NJ, Santamaría L & Schwenk K. Clonal diversity and structure within a population of the pondweed *Potamogeton pectinatus* foraged by Bewick's swans, 2137

Hansen AJ see Pitra C

Hansen MM see Bekkevold D

Hansen MM. Estimating the long-term effects of stocking domesticated trout into wild brown trout (*Salmo trutta*) populations: an approach using microsatellite DNA analysis of historical and contemporary samples, 1003

Hansen MM, Ruzzante DE, Nielsen EE Bekkevold D & Mensberg K-LD. Long-term effective population sizes, temporal stability of genetic composition and potential for local adaptation in anadromous brown trout (*Salmo trutta*) populations, 2523

Hansson B & Westerberg L. On the correlation between heterozygosity and fitness in natural populations, 2467

Hardy ME, Grady JM & Routman EJ. Intraspecific phylogeography of the slender madtom: the complex evolutionary history of the Central Highlands of the United States, 2393

Harr B see Schlötterer C

Haryanti see Benzie JAH

Haselkorn TS see Janzen FJ

Hasselquist D see Waldenström J

Hausser J see Lugon-Moulin N

Hawkins A see Garnier-Géré PH

Heath DD, Busch C, Kelly J & Atagi DY. Temporal change in genetic structure and effective population size in steelhead trout (*Oncorhynchus mykiss*), 197

Hebert PDN see Adamowicz SJ

Hedin M & Wood DA. Genealogical exclusivity in geographically proximate populations of *Hypochilus thorelli* Marx (Araneae, Hypochilidae) on the Cumberland Plateau of North America, 1575

Hedrick PW, Lee RN & Garrigan D. Major histocompatibility complex variation in red wolves: evidence for common ancestry with coyotes and balancing selection, 1905

Helbig AJ see Bensch S

Hellemans B see Hänfling B

Hellwig FH see Krüger AM

Henderson S see Trewick SA

Hendoux F see Bonnin I

Hendricks AC see Schultheis AS

Héral M see Garnier-Géré PH

Herre EA see Husband R

Hewitt GM see Ibrahim KM

Hewitt GM see Salomone N

Hoang TH see Huber K

Hoarau G, Rijnsdorp AD, Van Der Veer HW, Stam WT & Olsen JL. Population structure of plaice (*Pleuronectes platessa* L.) in northern Europe: microsatellites revealed large-scale spatial and temporal homogeneity, 1165

Hoegh-Guldberg O see Rodriguez-Lanetty M

Hoelzer G see McCafferty S

Högberg N, Kroken S, Thor G & Taylor JW. Reproductive mode and genetic variation suggest a North American origin of European *Letharia vulpina*, 1191

Holand Ø see Røed KH

Holder KR see Kimura M

Holderegger R see Stehlik I

Holderegger R, Stehlik I & Abbott RJ. Molecular analysis of the Pleistocene history of *Saxifraga oppositifolia* in the Alps, 1409

Holl A-C see Bonnin I

Holland BS & Hadfield MG. Islands within an island: phylogeography and conservation genetics of the endangered Hawaiian tree snail *Achatinella mustelina*, 365

Hollingsworth ML see Squirrell J

Hollingsworth PM see Squirrell J

Holsinger KE, Lewis PO & Dey DK. A Bayesian approach to inferring population structure from dominant markers, 1157

Hong D-Y see Song B-H

van Hooft WF, Groen AF & Prins HHT. Phylogeography of the African buffalo based on mitochondrial and Y-chromosomal loci: Pleistocene origin and population expansion of the Cape buffalo subspecies, 267

Hooper JNA see Wörheide G

Huang SSF, Hwang S-Y & Lin T-P. Spatial pattern of chloroplast DNA variation of *Cyclobalanopsis glauca* in Taiwan and East Asia, 2349

Huber K, Le Loan L, Hoang TH, Ravel S, Rodhain F & Failloux A-B. Genetic differentiation of the dengue vector, *Aedes aegypti* (Ho Chi Minh City, Vietnam) using microsatellite markers, 1629

Hurst GDD see Jiggins FM

Husband R see Vandenkoornhuyse P

Husband R, Herre EA, Turner SL, Gallery R & Young JPW. Molecular diversity of arbuscular mycorrhizal fungi and patterns of host association over time and space in a tropical forest, 2669

Huynen L, Millar CD & Lambert DM. A DNA test to sex ratite birds, 851

Hwang S-Y see Huang SSF

Ibrahim KM, Cooper SJB & Hewitt GM. Testing for recombination in a short nuclear DNA sequence of the European meadow grasshopper, Chorthippus parallelus, 583

Ijichi N see Kondo N

Imperatriz-Fonseca VL see Tóth E

Iraki N see Arafeh RMH

Irvine RJ see Côté SD

Irwin DE see Bensch S

Isagi Y see Kameyama Y

Jaarola M & Searle JB. Phylogeography of field voles (*Microtus agrestis*) in Eurasia inferred from mitochondrial DNA sequences, 2613

Jacobs DK see Dawson MN

Jameson R see Larson S

Jany J-L see Selosse M-A

Janzen FJ, Krenz JG, Haselkorn TS, Brodie Jr ED & Brodie III ED. Molecular phylogeography of common garter snakes (*Thamnophis sirtalis*) in western North America: implications for regional historical forces, 1739

Jarman SN, Gales NJ, Tierney M, Gill PC & Elliott NG. A DNA-based method for identification of krill species and its application to analysing the diet of marine vertebrate predators, 2679

Jarne P see Charbonnel N

Jarne P see Dutech C

Jarne P see Estoup A

Jarne P see Mavárez J

Jerome CA & Ford BA. The discovery of three genetic races of the dwarf mistletoe *Arceuthobium americanum* (Viscaceae) provides insight into the evoluiion of parasitic angiosperms, 387

Jerome CA & Ford BA. Comparative population structure and genetic diversity of *Arceuthobium americanum* (Viscaceae) and its *Pinus* host species: insight into host-parasite evolution in parasitic angiosperms, 407

Jesenšek D see Fumagalli L

Jiggins FM, Bentley JK, Majerus MEN & Hurst GDD. Recent changes in phenotype and patterns of host specialization in *Wolbachia* bacteria, 1275

Johnson EM see Freeman-Gallant CR

Johnson KP, Williams BL, Drown DM, Adams RJ & Clayton DH. The population genetics of host specificity: genetic differentiation in dove lice (Insecta: Phthiraptera), 25

Johnson NK & Cicero C. The role of ecologic diversification in sibling speciation of *Empidonax* flycatchers (Tyrannidae): multigene evidence from mtDNA, 2065

Johnson PCD, Whitfield JA, Foster WA & Amos W. Clonal mixing in the soldier-producing aphid *Pemphigus spyrothecae* (Hemiptera: Aphididae), 1525

Johnson WE see Palomares F

Joly HI see Dutech C

Jones PJ see Dallimer M

Jordan WC see Paulo OS

Joseph L, Wilke T & Alpers D. Reconciling genetic expectations from host specificity with historical population dynamics in an avian brood parasite, Horsfield's Bronze-Cuckoo *Chalcites basalis* of Australia, 829

Jug T see Fumagalli L

Kaderelt JW see Comes HP

Kai Y, Nakayama K & Nakabo T. Genetic differences among three colour morphotypes of the black rockfish, *Sebastes inermis*, inferred from mtDNA and AFLP analyses, 2591

Kalinowski ST. Evolutionary and statistical properties of three genetic distances, 1263

Kameyama Y, Isagi Y & Nakagoshi N. Relatedness structure in *Rhododendron metternichii* var. *hondoense* revealed by microsatellite analysis, 519

Kashani N see Dodd RS

Katoh H, Miura T, Maekawa K, Shinzato N & Matsumoto T. Genetic variation of symbiotic fungi cultivated by the macrotermitine termite *Odontotermes formosanus* (Isoptera: Termitidae) in the Ryukyu Archipelago, 1565

Katsoyannos BI see Bonizzoni M

Keim P see Mock KE

Kelly J see Heath DD

Kerdelhué C, Roux-Morabito G, Forichon J, Chambon J-M, Robert A & Lieutier F. Population genetic structure of *Tomicus piniperda* L. (Curculionidae: Scolytinae) on different pine species and validation of *T. destruens* (Woll.), 483

Kerth G, Mayer F & Petit E. Extreme sex-biased dispersal in the communally breeding, non-migratory Bechstein's bat (*Myotis bechsteinii*), 1491

Kiboi S see Waldenström J

Kiew R see Teo LL

Kimura M, Clegg SM, Lovette IJ, Holder KR, Girman DJ, Milá B, Wade P & Smith TB. Phylogeographical approaches to assessing demographic connectivity between breeding and overwintering regions in a Nearctic-Neotropical warbler (*Wilsonia pusilla*), 1605

King RA, Gornall RJ, Preston CD & Croft JM. Population differentiation of *Potamogeton pectinatus* in the Baltic Sea with reference to waterfowl dispersal , 1947

Kinnison MT, Bentzen P, Unwin MJ & Quinn TP. Reconstructing recent divergence: evaluating nonequilibrium population structure in New Zealand chinook salmon, 739

Kirschner J see Mes THM

Klopfenstein NB see Richardson BA

Knizhin I see Koskinen MT

Knop E see Ward PI

Knowles LL & Maddison WP. Statistical phylogeography, 2623

Knox D see Consuegra S

Koga R see Tsuchida T

Kondo N, Ijichi N, Shimada M & Fukatsu T. Prevailing triple infection with *Wolbachia* in *Callosobruchus chinensis* (Coleoptera: Bruchidae), 167

Korol AB see Li Y-C

Korpelainen H. A genetic method to resolve gender complements investigations on sex ratios in *Rumex acetosa*, 2151

Koskinen MT, Knizhin I, Primmer CR, Schlötterer C & Weiss S. Mitochondrial and nuclear DNA phylogeography of *Thymallus* spp. (grayling) provides evidence of ice-age mediated environmental perturbations in the world's oldest body of fresh water, Lake Baikal, 2599

Kosuch J see Vences M

Kotoulas G see Garnier-Géré PH

Kowalchuk GA, de Souza FA & van Veen JA. Community analysis of arbuscular mycorrhizal fungi associated with *Ammophila arenaria* in Dutch coastal sand dunes, 571

Kraus S see Waldick RC

Krebs CJ see Burton C

Kremer A see Mariette S

Kremer A see Ribeiro MM

Krenz JG see Janzen FJ

Krieg F see Triantafyllidis A

Kroken S see Högberg N

Kropf M see Comes HP

Krüger AM, Hellwig FH & Oberprieler C. Genetic diversity in natural and anthropogenic inland populations of salt-tolerant plants: random amplified polymorphic DNA analyses of *Aster tripolium* L. (Compositae) and *Salicornia ramosissima* Woods (Chenopodiaceae), 1647

Kuch M, Rohland N, Betancourt JL, Latorre C, Steppan S & Poinar HN. Molecular analysis of a 11 700-year-old rodent midden from the Atacama Desert, Chile, 913

Kuhn B see Sork VL

Kumpula J see Røed KH

Kuperus P see Mes THM

Kwon JA & Morden CW. Population genetic structure of two rare tree species (*Colubrina oppositifolia* and *Alphitonia ponderosa*, Rhamnaceae) from Hawaiian dry and mesic forests using random amplified polymorphic DNA markers, 991

Ladoukakis ED, Saavedra C, Magoulas A & Zouros E. Mitochondrial DNA variation in a species with two mitochondrial genomes: the case of *Mytilus galloprovincialis* from the Atlantic, the Mediterranean and the Black Sea, 755

Lambert DM see Huynen L

Langvatn R see Côté SD

Largiadèr CR see Gantenbein B

Larson S, Jameson R, Etnier M, Fleming M & Bentzen P. Loss of genetic diversity in sea otters (*Enhydra lutris*) associated with the fur trade of the 18th and 19th centuries, 1899

Latorre C see Kuch M

Law JH & Crespi BJ. The evolution of geographic parthenogenesis in *Timema* walking-sticks, 1471

Le Corre V see Mariette S

Le Loan L see Huber K

Leberg PL. Estimating allelic richness: Effects of sample size and bottlenecks, 2445

Lee RN see Hedrick PW

Lee RW see Cariello MO

Lee SK see Teo LL

Legg JP, French R, Rogan D, Okao-Okuja G & Brown JK. A distinct *Bemisia tabaci* (Gennadius) (Hemiptera: Sternorrhyncha: Aleyrodidae) genotype cluster is associated with the epidemic of severe cassava mosaic virus disease in Uganda, 1219

Leiss KA see Steinger T

Lemaire M see Vekemans X

Lenfant P see Planes S

Lepetit D, Brehm A, Fouillet P & Biémont C. Insertion polymorphism of retrotransposable elements in populations of the insular, endemic species *Drosophila madeirensis*, 347

Leterme N see Delmotte F

Lewis PO see Holsinger KE

Li Y-C, Korol AB, Fahima T, Beiles A & Nevo E. Microsatellites: genomic distribution, putative functions and mutational mechanisms: a review, 2453

Lieckfeldt D see Pitra C

Lieutier F see Kerdelhué C

Lilley JM see Moyes CL

Lin T-P see Huang SSF

Lindberg DR see Castilla JC

Lindell J see Primmer CR

Loeschcke V see Bekkevold D

Loeschcke V see Frydenberg J

Loriot S see Despres L

Louie KD see Dawson MN

Lovette IJ see Kimura M

Lübberstedt Th see Barth S

Lucchini V, Fabbri E, Marucco F, Ricci S, Boitani L & Randi E. Noninvasive molecular tracking of colonizing wolf (*Canis lupus*) packs in the western Italian Alps, 857

Lugon-Moulin N see Balloux F

Lugon-Moulin N & Hausser J. Phylogeographical structure, postglacial recolonization and barriers to gene flow in the distinctive Valais chromosome race of the common shrew (*Sorex araneus*), 785

Luikart G see Maudet C

Lumaret, R. Mir, C. Michaud H. & Raynal V. Phylogeographical variation of chloroplast DNA in holm oak (*Quercus ilex* L.), 2327

MacDonald C & Brookfield JFY. Intraspecific molecular variation in the seaweed fly *Coelopa frigida* consistent with behavioural distinctness of British and Swedish populations, 1637

Macedo RHF see Cariello MO

Mackiewicz M, Fletcher DE, Wilkins SD, DeWoody JA & Avise JC. A genetic assessment of parentage in a natural population of dollar sunfish (*Lepomis marginatus*) based on microsatellite markers, 1877

Maddison WP see Knowles LL

Maekawa K see Katoh H

Magoulas A see Garnier-Géré PH

Magoulas A see Ladoukakis ED

Majerus MEN see Jiggins FM

Major RE see Colgan D

Malacrida A see Bonizzoni M

Malhotra A see Stenson AG

Marguerie R see Bonizzoni M

Mariette S see Ribeiro MM

Mariette S, Le Corre V, Austerlitz F & Kremer A. Sampling within the genome for measuring within-population diversity: trade-offs between markers, 1145

Marinone MC see Adamowicz SJ

Márquez LM, van Oppen MJH, Willis BL, Reyes A & Miller DJ. The highly cross-fertile coral species, *Acropora hyacinthus* and *Acropora cytherea*, constitute statistically distinguishable lineages, 1339

Marshall F see Côté SD

Marshall HD & Ritland K. Genetic diversity and differentiation of Kermode bear populations, 685

Martinelli R see Saino N

Martínez Zapater JM see Arroyo García R

Marucco F see Lucchini V

Massonnet B, Simon J-C & Weisser WW. Metapopulation structure of the specialized herbivore *Macrosiphoniella tanacetaria* (Homoptera, Aphididae), 2511

Masutti L see Salvato P

Mathews LM, Schubart CD, Neigel JE & Felder DL. Genetic, ecological, and behavioural divergence between two sibling snapping shrimp species (Crustacea: Decapoda: *Alpheus*), 1427

Matocq MD. Phylogeographical structure and regional history of the dusky-footed woodrat, *Neotoma fuscipes*, 229

Matsumoto T see Katoh H

Matsumoto T see Tsuchida T

Matthee CA & Flemming AF. Population fragmentation in the southern rock agama, *Agama atra*: more evidence for vicariance in Southern Africa, 465

Maudet C, Miller C, Bassano B, Breitenmoser-Würsten C, Gauthier D, Obexer-Ruff G, Michallet J, Taberlet P & Luikart G. Microsatellite DNA and recent statistical methods in wildlife conservation management: applications in Alpine Ibex [*Capra ibex* (*ibex*)], 421

Mavárez J, Amarista M, Pointier J-P & Jarne P. Fine-scale population structure and dispersal in *Biomphalaria glabrata*, the intermediate snail host of *Schistosoma mansoni*, in Venezuela, 879

May-Itza WJ see Palmer KA

Mayer F see Kerth G

McCafferty S, Bermingham E, Quenouille B, Planes S, Hoelzer G & Asoh K. Historical biogeography and molecular systematics of the Indo-Pacific genus *Dascyllus* (Teleostei: Pomacentridae), 1377

McMillan WO see Rivera-Ocasio E

McPeek MA see Turgeon J

McPheron BA see Meixner MD

Meixner MD, McPheron BA, Silva JG, Gasparich GE & Sheppard WS. The Mediterranean fruit fly in California: evidence for

multiple introductions and persistent populations based on microsatellite and mitochondrial DNA variability, 891
Melchinger AE see Barth S
Menozzi P see Bucci G
Mensbery K-LD see Hansen MM
Mes THM, Kuperus P, Kirschner J, Štepánek J, Štorchová H, Oosterveld P & den Nijs JCM. Detection of genetically divergent clone mates in apomictic dandelions, 253
Meusnier I, Valero M, Destombe C, Godé C, Desmarais E, Bonhomme F, Stam WT & Olsen JL. Polymerase chain reaction–single strand conformation polymorphism analyses of nuclear and chloroplast DNA provide evidence for recombination, multiple introductions and nascent speciation in the *Caulerpa taxifolia* complex, 2317
Meyer A see Vences M
Meyer CP see Castilla JC
Michallet J see Maudet C
Michand H see Lumaret R
Milá B see Kimura M
Milinkovitch MC see Ciofi C
Milinkovitch MC see Fournier D
Millar CD see Huynen L
Miller C see Maudet C
Miller DJ see Márquez LM
Miller DJ see van Oppen MJH
Mir C see Lumaret R
Mitchell-Olds T see Clauss MJ
Miura T see Katoh H
Mizen S see Sackville Hamilton NR
Mizen S see Skøt L
Mock KE, Theimer TC, Rhodes Jr OE, Greenberg DL & Keim P. Genetic variation across the historical range of the wild turkey (*Meleagris gallopavo*), 643
Mokady O see Appelbaum L
Møller AP see Saino N
Moore MK & Ball Jr RM. Multiple paternity in loggerhead turtle (*Caretta caretta*) nests on Melbourne Beach, Florida: a microsatellite analysis, 281
Moosa MK see Barber PH
Moran GF see Burczyk J
Moran NA see Abbot P
Morand M-E, Brachet S, Rossignol P, Dufour J & Frascaria-Lacoste N. A generalized heterozygote deficiency assessed with microsatellites in French common ash populations, 377
Morand-Prieur ME, Vedel F, Raquin C, Brachet S, Sihachakr D & Frascaria-Lacoste N. Maternal inheritance of a chloroplast microsatellite marker in controlled hybrids between *Fraxinus excelsior* and *Fraxinus angustifolia*, 613
Morden CW see Kwon JA
Morgan-Richards M see Trewick SA
Moria S see Benzie JAH
Morrison SF, Scott Keogh J & Scott IAW. Molecular determination of paternity in a natural population of the multiply mating polygynous lizard *Eulamprus heatwolei*, 535
Moyes CL, Lilley JM, Casais CA, Cole SG, Haeger PD & Dale PJ. Barriers to gene flow from oilseed rape (*Brassica napus*) into populations of *Sinapis arvensis*, 103
Mueller UG see Green AM
Müller-Schärer H see Steinger T
Mundy NI see Surridge AK
Murray TD see Douhan GW
Naciri-Graven Y see Garnier-Géré PH
Nakabo T see Kai Y
Nakagoshi N see Kameyama Y
Nakayama K see Kai Y
Neigel JE see Mathews LM
Nevo E see Li Y-C
Nichols RA see Paulo OS
Nieminen M see Røed KH
Nielsen EE see Hansen MM
Nieto Feliner G see Gutiérrez Larena B
Nieves-Aldrey J-L see Rokas A
den Nijs JCM see Mes THM
Niklfeld H see Schönswetter P
Nogueira-Neto P see Tóth E

Oberprieler C see Krüger AM
Obexer-Ruff G see Maudet C
O'Brien SJ see Palomares F
O'Brien SJ see Comstock KE
O'Connor M see Crim JL
Ogden R & Thorpe RS. The usefulness of amplified fragment length polymorphism markers for taxon discrimination across graduated fine evolutionary levels in Caribbean *Anolis* lizards, 437
Okao-Okuja G see Legg JP
Oldroyd BP see Palmer KA
Olsen JL see Hoarau G
Olsen JL see Meusnier I
Olsen JL see Van der Strate HJ
Olsen KM. Population history of *Manihot esculenta* (Euphorbiaceae) inferred from nuclear DNA sequences, 901
Oosterveld P see Mes THM
van Oppen MJH see Márquez LM
van Oppen MJH see Shearer TL
van Oppen MJH, Willis BL, van Rheede T & Miller DJ. Spawning times, reproductive compatibilities and genetic structuring in the *Acropora aspera* group: evidence for natural hybridization and semi-permeable species boundaries in corals, 1363
Ostrander EA see Comstock KE
Ottosson U see Waldenström J
Ouborg NJ see Hangelbroek HH
Owens IPF see Griffith SC

Paladino FV see Crim JL
Palmé AE & Vendramin GG. Chloroplast DNA variation, postglacial recolonization and hybridization in hazel, *Corylus avellana*, 1769
Palmer KA, Oldroyd BP, Quezada-Euán JJG, Paxton RJ & May-Itza WJ. Paternity frequency and maternity of males in some stingless bee species, 2107
Palomares F, Godoy JA, Piriz A and O'Brien SJ & Johnson WE. Faecal genetic analysis to determine the presence and distribution of elusive carnivores: design and feasibility for the Iberian lynx, 2171
Palumbi SR see Barber PH
Papert A see Floyd R
Patarnell T see Salvato P
Patarnello T see Zane L
Paulo OS, Jordan WC, Bruford MW & Nichols RA. Using nested clade analysis to assess the history of colonization and the persistence of populations of an Iberian Lizard, 809
Paxton RJ see Palmer KA

Paxton RJ, Ayasse M, Field J & Soro A. Complex sociogenetic organization and reproductive skew in a primitively eusocial sweat bee, *Lasioglossum malachurum*, as revealed by microsatellites, 2405
Pearson CVM, Rogers AD & Sheader M. The genetic structure of the rare lagoonal sea anemone, *Nematostella vectensis* Stephenson (Cnidaria; Anthozoa) in the United Kingdom based on RAPD analysis, 2285
Pecchioli E see Vernesi C
Pecon-Slattery J see Comstock KE
Peeters C see Doums C
Peever TL see Douhan GW
Pelton MR see Ratnayeke S
Pemberton JM see Dallimer M
Peng P-H see Song B-H
Perrin N see Ehinger M
Perrin N see Goudet J
Persat H see Weiss S
Pertoldi C see Frydenberg J
Petit E see Ehinger M
Petit E see Kerth G
Petit RJ see Grivet D
Petronelli P see Dutech C
Pfenninger M see Wilke T
Pilastro A see Saino N
Pintér I see Trewick SA
Pinto J, Donnelly MJ, Sousa CA, Gil V, Ferreira C, Elissa N, Do Rosário VE & Charlwood JD. Genetic structure of *Anopheles gambiae* (Diptera: Culicidae) in São Tomé and Príncipe (West Africa): implications for malaria control, 2183
Piriz A see Palomares F
Pitra C, Hansen AJ, Lieckfeldt D & Arctander P. An exceptional case of historical outbreeding in African sable antelope populations, 1197
Planes S see McCafferty S
Planes S & Lenfant P. Temporal change in the genetic structure between and within cohorts of a marine fish, *Diplodus sargus*, induced by a large variance in individual reproductive success, 1515
Plomion C see Ribeiro MM
Poinar HN see Kuch M
Pointier J-P see Mavárez J
Porter BA see Fiumera AC
Porter BA see Walker D
Potti J, Dávila JA, Tella JL, Frías Ó & Villar S. Gender and viability selection on morphology in fledgling pied flycatchers, 1317
Powell JR see Ciofi C
Preston CD see King RA
Primmer CR see Koskinen MT
Primmer CR, Borge T, Lindell J & Sætre G-P. Single-nucleotide polymorphism characterization in species with limited available sequence information: high nucleotide diversity revealed in the avian genome, 603
Prins HHT see van Hooft WF
Prugnolle F, De Meeûs T, Durand P, Sire C & Théron A. Sex-specific genetic structure in *Schistosoma mansoni*: evolutionary and epidemiological implications, 1231

Queller DC see Tóth E
Quenouille B see McCafferty S
Quezada-Euán JJG see Palmer KA
Quinn TP see Kinnison MT

Randi E see Lucchini V
Randi E see Vernesi C
Raquin C see Morand-Prieur ME
Rasatavonjizay R see Charbonnel N
Ratkiewicz M see Brzosko E
Ratnayeke S, Tuskan GA & Pelton MR. Genetic relatedness and female spatial organization in a solitary carnivore, the raccoon, *Procyon lotor*, 1115
Ratnieks FLW see Châline N
Rauscher JT, Doyle JJ & Brown AHD. Internal transcribed spacer repeat-specific primers and the analysis of hybridization in the *Glycine tomentella* (Leguminosae) polyploid complex, 2691
Ravel S see Huber K
Raynal see Lumaret R
Reina R see Crim JL
Rendell S & Ennos RA. Chloroplast DNA diversity in *Calluna vulgaris* (heather) populations in Europe, 69
Reyes A see Márquez LM
van Rheede T see van Oppen MJH
Rhodes Jr OE see Mock KE
Rhodes Jr OE see Williams CL
Ribeiro MM, Mariette S, Vendramin GG, Szmidt AE, Plomion C & Kremer A. Comparison of genetic diversity estimates within and among populations of maritime pine using chloroplast simple-sequence repeat and amplified fragment length polymorphism data, 869
Ricci S see Lucchini V
Richardson BA, Brunsfeld SJ & Klopfenstein NB. DNA from bird-dispersed seed and wind-disseminated pollen provides insights into postglacial colonization and population genetic structure of whitebark pine (*Pinus albicaulis*), 215
Riço C & Turner GF. Extreme microallopatric divergence in a cichlid species from Lake Malawi, 1585
Riegler M & Stauffer C. *Wolbachia* infections and superinfections in cytoplasmically incompatible populations of the European cherry fruit fly *Rhagoletis cerasi* (Diptera, Tephritidae), 2425
Rieseberg LH see Schwarzbach AE
Rijnsdorp AD see Hoarau G
Rispe C see Delmotte F
Ritland K see Marshall HD
Rivera-Ocasio E, Aide TM & McMillan WO. Patterns of genetic diversity and biogeographical history of the tropical wetland tree, *Pterocarpus officinalis* (Jacq.), in the Caribbean Basin, 675
Robert A see Kerdelhué C
Robertson DR see Rocha LA
Roca AL see Comstock KE
Rocha LA, Bass AL, Robertson DR & Bowen BW. Adult habitat preferences, larval dispersal, and the comparative phylogeography of three Atlantic surgeonfishes (Teleostei: Acanthuridae), 243
Rodhain F see Huber K
Rodriguez-Lanetty M & Hoegh-Guldberg O. The phylogeography and connectivity of the latitudinally widespread scleractinian coral *Plesiastrea versipora* in the Western Pacific, 1177
Røed KH, Holand Ø, Smith ME, Gjøstein H, Kumpula J & Nieminen M. Reproductive success in reindeer males in a herd with varying sex ratio, 1239
Rogan D see Legg JP
Rogers AD see Pearson CVM
Rohland N see Kuch M

Rokas A, Atkinson RJ, Nieves-Aldrey J-L, West SA & Stone GN. The incidence and diversity of *Wolbachia* in gallwasps (Hymenoptera Cynipidae) on oak, 1815
Roldán-Ruiz I see Vekemans X
Romano SL see Shearer TL
Rossignol P see Morand M-E
Routman EJ see Hardy ME
Roux-Morabito G see Kerdelhué C
Rueness J see Gabrielsen TM
Rumsey FJ see Trewick SA
Russell SJ see Trewick SA
Ruzzante DE see Hansen MM
Ryan Gregory T see Adamowicz SJ

Saavedra C see Ladoukakis ED
Sackville Hamilton NR see Skøt L
Sackville Hamilton NR, Skøt L, Chorlton KH, Thomas ID & Mizen S. Molecular genecology of temperature response in *Lolium perenne*: 1. preliminary analysis to reduce false positives, 1855
Sætre G-P see Primmer CR
Saino N, Ambrosini R, Martinelli R, Calza S, Møller AP & Pilastro A. Offspring sexual dimorphism and sex-allocation in relation to parental age and paternal ornamentation in the barn swallow, 1533
Sait SM see Burden JP
Salomon M see Bensch S
Salomone N, Emerson BC, Hewitt GM & Bernini F. Phylogenetic relationships among the Canary Island Steganacaridae (Acari, Oribatida) inferred from mitochondrial DNA sequence data, 79
Salvato P, Battisti A, Concato S, Masutti L, Patarnell T & Zane L. Genetic differentiation in the winter pine processionary moth (*Thaumetopoea pityocampa — wilkinsoni* complex), inferred by AFLP and mitochondrial DNA markers, 2435
Salzburger W, Baric S & Sturmbauer C. Speciation via introgressive hybridization in East African cichlids?, 619
Santamaría L see Hangelbroek HH
Sapir Y see Arafeh RMH
Saponara F see Freeman-Gallant CR
Schaefer H-C see Vences M
Schliewen U see Englbrecht CC
Schlötterer C see Koskinen MT
Schlötterer C see Weiss S
Schlötterer C & Harr B. Single nucleotide polymorphisms derived from ancestral populations show no evidence for biased diversity estimates in *Drosophila melanogaster*, 947
Schneeder S see Dupamloup I
Scholte E-J see Ward PI
Schönswetter P, Tribsch A, Barfuss M and Niklfeld H. Several Pleistocene refugia detected in the high alpine plant *Phyteuma globulariifolium* Sternb. & Hoppe (Campanulaceae) in the European Alps, 2637
Schubart CD see Mathews LM
Schultheis AS, Weigt LA & Hendricks AC. Gene flow, dispersal, and nested clade analysis among populations of the stonefly *Peltoperla tarteri* in the southern Appalacians, 317
Schwabb HG see Cariello MO
Schwarzbach AE & Rieseberg LH. Likely multiple origins of a diploid hybrid sunflower species, 1703
Schwenk K see Hangelbroek HH
Scott IAW see Morrison SF
Scott Keogh J see Morrison SF
Scott Mills L see Tallmon DA
Searle JB see Jaarola M
Segelbacher G & Storch I. Capercaillie in the Alps: genetic evidence of metapopulation structure and population decline, 1669
Seibold I see Bensch S
Seiter J see Dutech C
Selosse M-A, Weiß M, Jany J-L & Tillier A. Communities and populations of sebacinoid basidiomycetes associated with the achlorophyllous orchid *Neottia nidus-avis* (L.) L.C.M. Rich. and neighbouring tree ectomycorrhizae, 1831
Serdio A see Consuegra S
Serfass TL see Williams CL
Sester M see Gueritaine G
Set O see Teo LL
Sheader M see Pearson CVM
Shearer TL, van Oppen MJH, Romano SL & Wörheide G. Slow mitochondrial DNA sequence evolution in the Anthozoa (Cnidaria), 2475
Sheppard WS see Meixner MD
Shibao H see Tsuchida T
Shimada M see Kondo N
Shinzato N see Katoh H
Shmida A see Arafeh RMH
Sihachakr D see Morand-Prieur ME
Silva JG see Meixner MD
Simon J-C see Delmotte F
Simon J-C see Massonnet B
Sire C see Prugnolle F
Skøt L see Sackville Hamilton NR
Skøt L, Sackville Hamilton NR, Mizen S, Chorlton KH & Thomas ID. Molecular genecology of temperature response in *Lolium perenne*: 2. association of AFLP markers with ecogeography, 1865
Slippers B, Wingfield BD, Coutinho TA & Wingfield MJ. DNA sequence and RFLP data reflect geographical spread and relationships of *Amylostereum areolatum* and its insect vectors, 1845
Smith ME see Røed KH
Smith P see Burden JP
Smith TB see Kimura M
Smouse PE see Sork VL
Snoj A see Fumagalli L
Song B-H, Wang X-Q, Wang X-R, Sun L-J, Hong D-Y & Peng P-H. Maternal lineages of *Pinus densata*, a diploid hybrid, 1057
Sork VL, Davis FW, Smouse PE, Apsit VJ, Dyer RJ, Fernandez-M JF & Kuhn B. Pollen movement in declining populations of California Valley oak, *Quercus lobata*: where have all the fathers gone?, 1657
Soro A see Paxton RJ
Sousa CA see Pinto J
de Souza FA see Kowalchuk GA
Spotila JR see Crim JL
Spotila LD see Crim JL
Squirrell J, Hollingsworth PM, Bateman RM, Tebbitt MC & Hollingsworth ML. Taxonomic complexity and breeding system transitions: conservation genetics of the *Epipactis leptochila* complex (Orchidaceae), 1957
Stabile J see Grunwald C
Stam WT see Hoarau G
Stam WT see Meusnier I
Stam WT see Van der Strate HJ

Stanger M see Freeman-Gallant CR
Stauffer C see Riegler M
Stehlik I see Holderegger R
Stehlik I, Blattner FR, Holderegger R & Bachmann K. Nunatak survival of the high Alpine plant *Eritrichium nanum* (L.) Gaudin in the central Alps during the ice ages, 2027
Steinger T, Haldimann P, Leiss KA & Müller-Schärer H. Does natural selection promote population divergence? A comparative analysis of population structure using amplified fragment length polymorphism markers and quantitative traits, 2583
Stenson AG, Malhotra A & Thorpe RS. Population differentiation and nuclear gene flow in the Dominican anole (*Anolis oculatus*), 1679
Štepánek J see Mes THM
Steppan S see Kuch M
Stone GN see Rokas A
Stone KD, Flynn RW & Cook JA. Post-glacial colonization of northwestern North America by the forest-associated American marten (*Martes americana*, Mammalia: Carnivora: Mustelidae), 2049
Storch I see Segelbacher G
Štorchová H see Mes THM
Storz JF. Contrasting patterns of divergence in quantitative traits and neutral DNA markers: analysis of clinal variation, 2537
Strassmann JE see Tóth E
Straus LG see Consuegra S
Sturmbauer C see Salzburger W
Sugama K see Benzie JAH
Sun L-J see Song B-H
Surridge AK & Mundy NI. Trans-specific evolution of opsin alleles and the maintenance of trichromatic colour vision in Callitrichine primates, 2157
Swift CC see Dawson MN
Symondson WOC. Molecular identification of prey in predator diets, 627
Szmidt AE see Ribeiro MM

Taberlet P see Maudet C
Tallmon DA, Draheim HM, Scott Mills L & Allendorf FW. Insights into recently fragmented vole populations from combined genetic and demographic data, 699
Tautz D see Englbrecht CC
Taylor EB see Burton C
Taylor JW see Högberg N
Tebbitt MC see Squirrell J
Tella JL see Potti J
Teo LL, Kiew R, Set O, Lee SK & Gan YY. Hybrid status of kuwini, *Mangifera odorata* Griff. (Anacardiaceae) verified by amplified fragment length polymorphism, 1465
Theimer TC see Mock KE
Théron A see Prugnolle F
Thomas ID see Sackville Hamilton NR
Thomas ID see Skøt L
Thor G see Högberg N
Thorpe RS see Ogden R
Thorpe RS see Stenson AG
Thuman, KA see Griffith SC
Tiedemann R see Vernesi C
Tierney M see Jarman SN
Tillier A see Selosse M-A
Tóth E, Strassmann JE, Nogueira-Neto P, Imperatriz-Fonseca VL & Queller DC. Male production in stingless bees: variable outcomes of queen–worker conflict, 2661
Trewick SA, Morgan-Richards M, Russell SJ, Henderson S, Rumsey FJ, Pintér I, Barrett JA, Gibby M & Vogel JC. Polyploidy, phylogeography and Pleistocene refugia of the rockfern *Asplenium ceterach*: evidence from chloroplast DNA, 2003
Triantafyllidis A, Krieg F, Cottin C, Abatzopoulos TJ, Triantaphyllidis C & Guyomard R. Genetic structure and phylogeography of European catfish (*Silurus glanis*) populations, 1039
Triantaphyllidis C see Triantafyllidis A
Tribsch A see Schönswetter P
Tsuchida T, Koga R, Shibao H, Matsumoto T & Fukatsu T. Diversity and geographic distribution of secondary endosymbiotic bacteria in natural populations of the pea aphid, *Acyrthosiphon pisum*, 2123
Turgeon J & McPeek MA. Phylogeographic analysis of a recent radiation of *Enallagma* damselflies (Odonata: Coenagrionidae), 1989
Turner GF see Riço C
Turner SL see Husband R
Tuskan GA see Ratnayeke S

Uiblein F see Weiss S
Unwin MJ see Kinnison MT
Usher AV see England PR

Valero M see Meusnier I
Väli Ü. Mitochondrial pseudo-control region in old world eagles (genus *Aquila*), 2189
Valsecchi E, Hale P, Corkeron P & Amos W. Social structure in migrating humpback whales (*Megaptera novaeangliae*), 507
Van de Zande L see Van der Strate HJ
Van der Strate HJ, Van de Zande L, Stam WT & Olsen JL. The contribution of haploids, diploids and clones to fine-scale population structure in the seaweed *Cladophoropsis membranacea* (Chlorophyta), 329
Van Der Veer HW see Hoarau G
Vandenkoornhuyse P, Husband R, Daniell TJ, Watson IJ, Duck JM, Fitter AH & Young JPW. Arbuscular mycorrhizal community composition associated with two plant species in a grassland ecosystem, 1555
Vedel F see Morand-Prieur ME
van Veen JA see Kowalchuk GA
Veith M see Vences M
Vekemans X, Beauwens T, Lemaire M & Roldán-Ruiz I. Data from amplified fragment length polymorphism (AFLP) markers show indication of size homoplasy and of a relationship between degree of homoplasy and fragment size, 139
Vences M, Andreone F, Glaw F, Kosuch J, Meyer A, Schaefer H-C & Veith M. Exploring the potential of life-history key innovation: brook breeding in the radiation of the Malagasy treefrog genus *Boophis*, 1453
Vendramin GG see Palmé AE
Vendramin GG see Ribeiro MM
Vernesi C, Pecchioli E, Caramelli D, Tiedemann R, Randi E & Bertorelle G. The genetic structure of natural and reintroduced roe deer (*Capreolus capreolus*) populations in the Alps and central Italy, with reference to the mitochondrial DNA phylogeography of Europe, 1285

Verspoor E see Consuegre S
Viard F see Bonnin I
Villar S see Potti J
Vitalis R. Sex-specific genetic differentiation and coalescence times: estimating sex-biased dispersal rates, 125
Vogel JC see Trewick SA
Volckaert FAM see Hänfling B
Vonwil J see Ward PI

Wade P see Kimura M
Waldenström J, Bensch S, Kiboi S, Hasselquist D & Ottosson U. Cross-species infection of blood parasites between resident and migratory songbirds in Africa, 1545
Waldick RC, Kraus S, Brown M & White BN. Evaluating the effects of historic bottleneck events: an assessment of microsatellite variability in the endangered, North Atlantic right whale, 2241
Waldman JR see Grunwald C
Walker D, Porter BA & Avise JC. Genetic parentage assessment in the crayfish *Orconectes placidus*, a high-fecundity invertebrate with extended maternal brood care, 2115
Wang X-Q see Song B-H
Wang X-R see Song B-H
Waples RS. Evaluating the effect of stage-specific survivorship on the N_e/N ratio, 1029
Ward PI, Vonwil J, Scholte E-J & Knop E. Field experiments on the distribution of eggs of different phosphoglucomutase (PGM) genotypes in the yellow dung fly *Scathophaga stercoraria* (L.), 1781
Wares JP. Community genetics in the Northwestern Atlantic intertidal, 1131
Waser P see Goudet J
Wasser SK see Comstock KE
Watson IJ see Vandenkoornhuyse P
Weiblen GD & Bush GL. Speciation in fig pollinators and parasites, 1573
Weigt LA see Schultheis AS
Weiß M see Selosse M-A
Weiss S see Koskinen MT
Weiss S, Persat H, Eppe R, Schlötterer C & Uiblein F. Complex patterns of colonization and refugia revealed for European grayling *Thymallus thymallus*, based on complete sequencing of the mitochondrial DNA control region, 1393
Weisser WW see Massonnet B
West SA see Rokas A
Westerberg L see Hansson B
Whelan RJ see England PR
White BN see Waldick RC
Whitfield JA see Johnson PCD
Wilke T & Pfenninger M. Separating historic events from recurrent processes in cryptic species: phylogeography of mud snails (*Hydrobia* spp.), 1439
Wilke T see Joseph L
Wilkins SD see Mackiewicz M
Williams BL see Johnson KP
Williams CJ see Crim JL
Williams CL, Serfass TL, Cogan R & Rhodes Jr OE. Microsatellite variation in the reintroduced Pennsylvania elk herd, 1299
Willis BL see Márquez LM
Willis BL see van Oppen MJH
Wingfield BD see Slippers B
Wingfield MJ see Slippers B
Wirgin I see Grunwald C
Wood DA see Hedin M
Wörheide G see Shearer TL
Wörheide G, Hooper JNA & Degnan BM. Phylogeography of western Pacific *Leucetta 'chagosensis'* (Porifera: Calcarea) from ribosomal DNA sequences: implications for population history and conservation of the Great Barrier Reef World Heritage Area (Australia), 1753
Wright C see Cowgill SE
Wróblewska A see Brzosko E

Young JPW see Husband R
Young JPW see Vandenkoornhuyse P

Zabalgogeazcoa I see Arroyo García R
Zane L see Salvato P
Zane L, Bargelloni L & Patarnello T. Strategies for microsatellite isolation: a review, 1
Zouros E see Ladoukakis ED

Molecular Ecology (2002) **11**, 2729–2734

Keyword index

16S mtDNA sequences, 925
16S rRNA, 55, 465
18S ribosomal RNA (SSU), 839
412, 347

Acanthurus, 243
Achatinella, 365
Acipenseriformes, 1885
Acraea, 1275
Acropora, 1363
Acyrthosiphon pisum, 2123
adaptive differentiation, 1679
adaptive variation, 1905
admixture, 1197, 2213
admixture analysis, 1003
Aedes aegypti, 1629
AFLP, 139, 355, 387, 407, 437, 473, 643, 675, 869, 1327, 1465, 2229, 2359, 2591
AFLP markers, 1145
African elephant, 2489
Agamidae, 465
Agaonidae, 1573
aldicarb, 821
alga, 329
alleles per locus, 2445
allelic diversity, 2445
allo polyploid, 1351
allopatric divergence, 2065
allopatry, 1439
allozyme, 113, 711, 1515
Allozyme diversity, 1499
allozymes, 925, 1781, 1957, 2499
Alpheus, 1427
Alpine species, 1409
Alps, 1669
altruism, 1525
AM1, 2669
American martens, 2049
AMF, 571
AMOVA, 483, 2571
Amphibia, 1453
amplified fragment length polymorphism, 2637
amplified fragment length polymorphism (AFLP), 1157
amplified fragment length polymorphism markers (AFLP), 2337
amplified fragment length polymorphisms, 447
anarchy, 1795
ancient DNA, 913, 1899
ancient mtDNA, 2037
Anolis, 437
Anopheles gambiae, 2183
anthozoa, 2475
Anthyllis montana, 447
ants, 1805
AP-PCR, 1565
aphid, 2123, 2511
aphids, 2649
Apis mellifera, 1795
Arabidopsis lyrata, 591
Arabidopsis thaliana, 495, 591
arbuscular mycorrhiza, 2669
arbuscular mycorrhizal (AM) fungi, 1555
Arceuthobium americanum, 387, 407
area cladogram, 2393
Armeria, 1965
asexual reproduction, 2083, 2317
asexuality, 711
assignment index, 529, 1103
assignment tests, 289, 421
associative overdominance, 2467
Aster tripolium, 1647
Atacama Desert, 913
Atlantic salmon, 2037
Attini, 191
Austroplebeia, 2107
avian breeding systems, 2195
Avicennia germinans, 1327

B-complex, 1125
backcrosses, 1419
Baltic Sea, 2083
barnacles, 113
barrier, 2349
Basidiomycetes, 1845
bats, 2537
Bayesian modelling, 1157
Bergmann's rule, 2537
bias, 947
bilbo, 347
biodiversity, 1957
biodiversity assessment, 437, 839
biogeography, 215, 243, 1191, 1327, 1637, 2553, 2623
Biomphalaria glabrata, 879
biosafety, 821
bird, 603, 2189
birds, 851
Boechera, 591
bottleneck, 591, 1923, 2213, 2445
bottleneck detection, 421
Brassica napus, 103
Brazil, 243
breeding dispersal, 289
breeding system, 1931
brood parasites, 829
brood parasitism, 1877
Buchnera, 2649

Caesalpiniaceae, 725
California, 229
California Valley oak, 1657
Callitrichine primates, 2157
Callosobruchus chinensis, 167
Calluna vulgaris, 69
Canary Islands, 79
Canis lupus, 857
capercaillie, 1669
Caproleus caproleus, 2571
CAPS, 495
Caretta caretta, 281
Caribbean, 243
caribou, 1239
cassava mosaic geminiviruses, 1219
cDNA microarray, 17
central Atlantic islands, 243
Ceramium tenuicorne, 2083
Ceratitis capitata, 891
Cervus elaphus, 1299
cetaceans, 507
Chalcites, 829
charlock, 103
CHD1 genes, 1317
Chile, 1579
Chiroptera, 1491
chloroplast 16S intron, 2317
chloroplast DNA, 253, 613, 1351, 1769, 1965, 2003
chloroplast DNA (cpDNA), 69
chloroplast microsatellites, 69, 215, 869
cichlids, 619, 1585
Cirripedia, 1731
climate change, 913
clonal lineages, 355
clonal mixing, 1525
clonal reproduction, 1191, 1845
clonality, 329, 2499
clone mates, 253
CMD, 1219
coalescence, 2623
coalescence times, 125
coalescent, 1439
coalescent simulations, 2571
coastal marine evolution, 1065
Coelopa frigida, 1637
coevolution, 191, 557, 1573
cohort, 1515
COI gene, 79
cold tolerance, 1855, 1865
colonization, 951, 1017, 2049, 2213, 2511
colonization history, 181, 2613

© 2002 Blackwell Science Ltd

colour morphotypes, 2591
colour vision, 2157
columbiformes, 25
common ash, 377
communal breeding, 2417
community composition, 1555, 2669
community genetics, 1131
computer simulations, 771
concerted evolution, 2691
connectivity, 1605
conservation, 2703
conservation genetics, 39, 365, 421, 685, 1197, 2711
contact zone, 447
contemporary evolution, 739
control region, 229, 643, 1197, 1393, 1885
control-region, 1017
controlled crosses, 613
convergence, 1731
coral, 2475
corals, 1363
correlation, 1145
Corylus avellana, 1769
cospeciation, 1573
cost of migration, 1545
cost of sons and daughters, 1533
Cottidae, 1717
cpDNA, 979, 1957, 2027
cpDNA RFLP variation, 2327
cpDNA RFLPs, 1409
cpSSR, 869
Crassostrea gigas, 1499
cryptic female choice, 1781
cuckoos, 829
cyclical parthenogenesis, 711
Cynopterus sphinx, 2537
Cyphomyrmex, 191
Cypripedium calceolus, 2499
cystatin, 821
cytochrome b, 795
cytochrome *b*, 229, 465, 1545, 2049, 2171, 2613
cytochrome oxidase I, 25
cytoplasmic incompatibility, 167, 1275, 2425

D-loop, 267, 2171
Daktulosphaira vitifoliae, 2013
damselfish, 1377
de novo mutations, 2115
dengue, 1629
Dermochelys coriacea, 2097
Diacamma cyaneiventre, 2251
diet, 2679
differentiation, 2349
dioecy, 2151
Diptera, 1781
dispersal, 243, 317, 329, 659, 879, 1065, 1491, 1525, 1669, 1689, 1931
dispersal distances, 289
divergence, 253
diversification, 951
diversity, 947, 1351, 1555, 2669
DNA, 2649
DNA content, 1209
DNA microsatellites, 1795
DNA sequence analyses, 583
DNA sequence, 839, 947, 2003
Dodecanese, 925
dominance hierarchy, 1239
dominant markers, 1157
Doubly Uniparental Inheritance, 755
drift, 1515, 2583
Drosophila, 947, 1311
Drosophila buzzatii, 181
Drosophila madeirensis, 347
dung, 267
dynamics, 1555

EACMV-Ug, 1219
ecogeography, 1865
ecological divergence, 1427
ecology, 951
ectomycorrhiza, 1831
ectoparasites, 25
effective number of breeders, 2367
effective population size, 91, 739, 1029, 2241, 2523
effective size, 1263, 1491
Egernia stokesii, 1787
egg ejection, 2417
egg maternity, 2417
Electromorph, 1591
electrophoresis, 627
Empidonax, 2065
Enallagma, 1989
endangered plant, 2499
endangered plants, 991
endangered species, 1885, 2189
endemism, 795
endophytes, 355
endosymbionts evolution, 2649
Enhydra lutris, 1899
enrichment, 1
Epichloë festucae, 355
Ericaceae, 69
estimation, 1263
estuary, 1065
ESU, 365
Eubalaena australis, 2241
Eubalaena glacialis, 2241
Eurasia, 2613
Europe, 1393, 1717
European Alps, 2637
evergreen Mediterranean oaks, 2327
evolution, 465, 583, 951, 1209, 2475
evolution of sex, 711
Extinction, 2511
extra-pair paternity, 2195

F_{ST}, 771
F-statistics, 125, 155, 1103, 1157
Fabaceae, 447
faecal DNA, 2171
faeces, 2679
false positives, 1855
family correlated survival, 1029
family size variance, 1029
family structure, 1787
female dispersion, 1115
female remating, 1915
fennel pondweed, 2137
Festuca rubra, 355
field vole (*Microtus agrestis*), 2613
fitness, 1419, 2467
flatfish, 1165
flycatcher, 603
food web, 2679
Forest buffalo, 267
fossil middens, 913
founder events, 675
fragmentation, 967
Fraxinus excelsior, 377
Fraxinus species, 613
freshwater snail, 2213
ftsZ, 1815
functional overdominance, 2467
fungal population, 1831
fungi, 2229
fungus-growing ants, 191
fungus-growing termite, 1565

Gadus morhua, 91
Galápagos, 2265
gallwasp, 1815
gamergate, 2251
gametic disequilibrium, 2229
gene, 603
gene conversion, 583, 2157
gene diversity, 869, 1327
gene expression profiling, 17
gene flow, 55, 113, 215, 317, 519, 725, 739, 771, 979, 1177, 1419, 1637, 1689, 1931, 2265, 2307, 2379, 2435, 2523
gene-flow, 1947
genealogy, 2623
genecology, 1855, 1865
genetic analyses, 2171
genetic barriers, 2571
genetic differentiation, 25, 155, 437, 685, 1327, 1689, 1923, 2511
genetic distance, 1263
Genetic diversity, 1285
genetic diversity, 139, 197, 495, 1145, 1647, 1865, 1899, 1947, 2137, 2499
genetic drift, 939, 1689, 1923
genetic equilibrium, 739
genetic hotspot areas, 2027
genetic isolation, 1657
genetic markers, 305
genetic resources, 1855
genetic sex marker, 2151
genetic structure, 387, 407, 519, 675, 967, 1039, 1165, 2285, 2435, 2583

genetic variation, 377, 699, 1565, 2137, 2337
genetics, 659
genome heterogeneity, 1145
genomic distribution, 2453
genomics, 603
Geochelone, 2265
geographical distribution, 495, 2123
geographical space, 1115
glacial refugia, 795, 1393, 2613, 2637
glacial refugium, 2037
glacial survival, 2027
glaciation, 1285, 2599
glaciations, 1965
Glomeromycota, 571
Glomus, 571
glufosinate resistance, 1419
Glycine, 2691
GMO, 2703
granulin, 547
granulovirus, 547
grasslands, 355
Grevillea macleayana, 967
growth, 1499
guira cuckoo, 2417
Guira guira, 2417
gut analysis, 627
gypsy, 347

habitat fragmentation, 699, 1669, 2295, 2337
habitat shift, 2065
Haemoproteus, 1545
Halictidae, 2405
haplotype, 2349
Hawai'i, 991
Hawaiian tree snails, 365
Helianthus, 1703
heteroplasmy, 1885
heterozygosity, 591, 947, 1905, 1923, 2241, 2467
heterozygote deficiency, 377
hierarchical genetic structure, 2251
Hippotragus niger, 1197
Hirundo rustica, 1533
historical events, 939
historical samples, 2523
homoplasy, 139, 1591
horizontal transmission, 191, 1565, 1815
host-plant preference, 1555
host–parasite interactions, 1231
host shift, 1545
host specialization, 483
host specificity, 829
hybrid origin, 1465
hybrid speciation, 1703
hybrid zone, 473, 2359
hybridization, 55, 103, 925, 1057, 1209, 1339, 1363, 1703, 1769, 1965, 2591, 2691, 2711
Hydrobia, 1439
hydrography, 1617
Hypochilus, 1975

Iberian lynx, 2171
Iberian Peninsula, 809
IGS, 1845
immigration-selection balance, 1003
immunocompetence, 1125
inbreeding, 1491, 2467
incomplete lineage sorting, 2591
individual-based models, 1103
individual molecular tracking, 857
Indo-Pacific, 2553
Indo-West Pacific, 1377
inter al transcribed spacers, 447, 1753
introduction, 1579
introgression, 925, 1003, 1339, 1351, 1363, 2711
introgressive hybridization, 619
intron, 603
invasions, 2013
invasive species, 2317
inverse PCR, 2359
invertebrate, 2115
Iris, 39
island model, 125
island populations, 2499
islands, 951, 2183
isolation, 1, 1263
isolation-by-distance, 1805
isolation by distance, 289, 305, 1689, 2537
isolation of microsatellites, 181
isomorphic diplohaplontic life history, 329
isozymes, 979, 1781
ISSR, 495
ISSRs, 1947
ITS, 1831, 2691
ITS1, 483

join-count statistics, 305

Kermode bear, 685
kin selection, 2307
kinship, 507

Lacerta schreiberi, 809
lagoon, 2285
Lake Baikal, 2599
Lake Malawi, 1585
Lake Tanganyika, 619
landscape barriers, 1689
larvae, 659
latency, 547
laying order, 1533
leatherback, 2097
Lepidoptera, 1275, 2435
Leucetta 'chagosensis', 1753
lice, 25
lichenized fungi, 1191
life histories, 2151
life history evolution, 1377
likelihood method, 2661
linkage disequilibrium, 1905, 2467
lizard, 465
lizards, 1787
local adaptation, 2523
local genetic distinctiveness, 2295
loggerhead sea turtle, 281
Lolium perenne, 139
long-distance dispersal, 1191
Lycosidae, 2295

macroevolution, 1039
Madagascar, 1453
malaria control, 2183
male-killing, 1275
male parentage, 2661
Mangifera foetida, 1465
Mangifera indica, 1465
Mangifera odorata, 1465
mangrove, 1327
Manihot esculenta ssp. *flabellifolia*, 901
Manihot pruinosa, 901
Mantel test, 519
Mantellidae, 1453
marine fish, 1515
marine survival, 1029
marine, 659, 1947
mark–recapture, 2367
Martes, 2049
maternal inheritance, 613
maternity, 1877, 2115, 2367
mating system, 91, 535, 967, 1787, 2097, 2115, 2307, 2379
mating systems, 281
mating tactics, 1877
matriline, 2405
Mc1r, 685
mean-d^2, 1499
Mediterranean area, 2435
Mediterannean fruit fly, 1915
Meleagris gallopavo, 643
Meliponinae, 2661
metabolism, 939
metapopulation, 939, 1669, 2213, 2511
Mhc, 1125
microsatellite data, 1805
microsatellite DNA, 197, 289, 535, 1003, 2523, 2537
microsatellite loci, 619, 1679
microsatellite markers, 591
microsatellite, 113, 267, 281, 519, 529, 643, 711, 1165, 1499, 1923, 2097, 2511
microsatellites, 1, 155, 329, 377, 473, 507, 613, 685, 771, 785, 857, 879, 891, 967, 1017, 1039, 1103, 1145, 1231, 1299, 1311, 1491, 1525, 1585, 1591, 1629, 1669, 1787, 1899, 2115, 2183, 2241, 2251, 2265, 2359, 2489, 2661, 2711
migration, 507, 1263, 1605, 2307, 2349
migratory divide, 2359

mitochondrial and nuclear DNA sequences, 619
mitochondrial DNA, 79, 215, 229, 473, 795, 1057, 1197, 1377, 1545, 1637, 1885, 2065, 2295, 2475
mitochondrial DNA markers, 2171
mitotype, 1057
molecular divergence, 365
molecular markers, 1219, 1855, 1865, 2583
molecular phylogeny, 2065
molecular phylogeography, 1739
molecular sexing, 857, 1317
molecular systematics, 1377
molecular variance, 1647
monoclonal antibodies, 627
monogamy, 1787, 2195
Monotropa, 557
Monte Carlo, 1131
morphological convergence, 2327
morphological stasis, 1975
morphometric traits, 39
morphostasis, 1439
morphotypes, 79
*mt*CO1, 1219
mtDNA, 483, 829, 891, 1285, 1393, 1427, 1439, 1579, 1605, 1617, 1989, 2189, 2251
mtDNA control region, 317, 1065, 2591
mtDNA control-region, 857
mtDNA cox2–3 spacer, 2083
mtDNA genealogy, 1077
mtDNA sequences, 2013
mtDNA variation, 755, 2393, 2553
multiple-copy genes, 627
multiple origins, 2003
multiple paternity, 535, 2097, 2115
mussel, 1617
mussels, 755
Mustelidae, 795
mutation, 253, 1263, 2349
mutation models, 1591
mutualism, 1573
mycorrhizae, 557
Myotis bechsteinii, 1491
myxomatosis, 2703

*nad*1 intron, 1057
natural selection, 1317, 1679
ND4, 2393
neighbour-joining phylogeny, 197
nematode resistance, 821
nematodes, 839
Neotoma, 229
Neottia nidus-avis, 1831
nested clade analysis, 317, 901, 1077, 1177, 1197, 1339, 1471, 1753, 1989
nested cladistic analysis, 1975
networks, 2013
neutral theory, 1245
neutrality test, 1245
New Zealand, 1617
noninvasive, 2679
noninvasive genetics, 857
nontarget organisms, 821
northern refugia, 1717
Northwest Atlantic, 1131
Norway spruce, 305
nuclear cytoplasmic dis-equilibrium, 2317
nucleotide sequencing, 2393
nunataks, 2027

Odontotermes, 1565
oilseed rape, 103
opsin, 2157
orchids, 1831
oribatid mites, 79
Oryctolagus cuniculus, 2703
ostracods, 1931
outbreeding, 1197

Pacific Northwest, 2049
Pacific salmon, 1029
pairwise relatedness, 519
palaeohydrology, 2599
parasite transmission, 1545
parasitic angiosperm, 387, 407
parasitism, 17, 1317, 1573
parentage analysis, 2367, 2379
parental age, 1533
parthenogenesis, 1209, 1471
Partula, 55
paternity, 281, 1239, 1877, 2367
paternity assignment, 91
paternity frequency, 2107
PCR, 181, 2679
PCR-DGGE, 571
PCR-RFLP, 69, 891, 1957, 2027
PCR-RFLP fingerprinting, 1845
Pemphigus, 1525
Penaeus monodon, 2553
periglacial survival, 447
Perna canaliculus, 1617
Phaseolus lunatus, 139
phenotypic divergence, 2537
phenotypic plasticity, 1731, 2583
philopatry, 1115
Phylloscopus collybita, 473
Phylloscopus trochilus, 2359
Phyllotis, 913
phylogenetic tree, 1905
phylogeny, 951, 1453, 1573, 2189
phylogeographic structure, 365
phylogeography, 79, 229, 317, 465, 583, 643, 675, 809, 901, 913, 1039, 1065, 1131, 1177, 1377, 1393, 1409, 1439, 1471, 1605, 1617, 1703, 1717, 1753, 1769, 1965, 1975, 1989, 2037, 2049, 2327, 2393, 2489, 2599, 2613, 2623, 2637
pine forest, 2435
pink salmon, 1077
Pinus, 483, 1057
Pinus albi-caulis, 215
Pinus banksiana, 387, 407
Pinus contorta, 387, 407
Pinus pinaster, 869
plaice, 1165
plants, 967
Plasmodium, 1545
Plecoptera, 317
Pleistocene, 809, 901, 1717, 2003
Pleistocene glacial survival, 1409
Pleistocene glaciations, 1077
Plesiastrea versipora, 1177
Pleuronectes platessa, 1165
Pliocene, 809
pollen dispersal, 2379
pollen flow, 1657
pollen/seed migration, 979
polyandry, 281, 1805, 2097, 2405
polyclonal antisera, 627
polygyny, 1299, 1805, 2097, 2195
polymerase slippage, 2453
polymorphic allozyme, 2307
polymorphism, 1781, 2649
polyploidy, 2003, 2691
population bottleneck, 1899, 2241
population bottlenecks, 1299
population differentiation, 197, 869, 2337, 2583
population discrimination, 437
population divergence, 1285
population diversity, 2189
population dynamics, 699
population expansion, 181, 267
population genetic structure, 329, 1177
population genetics, 125, 591, 785, 991, 1191, 1299, 1591, 1617, 2229, 2265, 2623
population history, 39, 829
population structure, 17, 55, 113, 519, 771, 879, 1103, 1585, 1629, 1931, 1975, 2137, 2183, 2553, 2711
population structure and history, 1077
population structuring, 155, 347
population subdivision, 1311
population viscosity, 2251
populations, 2123
Porifera, 1753
postglacial colonization, 215
postglacial history, 69
postglacial recolonisation, 1351, 2337
postglacial recolonization, 1769
Potamogetonaceae, 1947
potsglacial evolution, 2037
prawn, 2553
presence and distribution, 2171
primary divergence, 39
principal component analysis, 785
propagule predation, 2137
Pteridophyte, 2003
Pyura praeputialis, 1579

Q_{ST}, 2537
Q. ilex L., 2327
quantitative genomics, 17
quantitative PCR, 167
Quaternary biogeography, 447, 2637
Quercus lobata, 1657

R_{ST}, 771
R-statistics, 155
rabbit haemorrhagic disease, 2703
radiation, 1989
random amplified polymorphic DNA (RAPD), 1157
randomization, 2445
randomizations, 1103
range expansion, 809, 829, 939
Rangifer, 1923
RAPD, 851, 991, 1115, 1231, 1647, 2083, 2137, 2285
RAPDs, 39, 1409
rarefaction, 2445
ratites, 851
Rattus rattus, 1231
rDNA, 1753
rDNA ITS, 1363
rDNA ITSs, 2317
recent long-distance dispersal, 2637
reciprocal monophyly, 1975
recolonization process, 725
recombination, 583, 1191, 2229, 2453
recurrent speciation, 1703
red-billed quelea, 529
reef fishes, 243
refugia, 1769
reindeer, 1239
reintroduction, 421, 1285
relatedness, 507
relative rate, 2475
relocation programs 1299
reproductive conflict, 2107
reproductive isolation, 1427, 2591
reproductive skew, 91
reproductive success, 1239
reproductive system, 711, 979
reptiles, 2265
reserves, 659
reticulate evolution, 1363
review, 1
RFLP, 613
Rfp-Y, 1125
Rhagoletis cerasi, 2425
Rhamnaceae, 991
Rhizoctonia, 1831
ribosomal DNA, 2691
risk assessment, 103, 821
roe deer, 1285
rRNA operon, 1845
RT–PCR, 547
Rumex acetosa, 2151
ryegrass, 1855, 1865

Salicornia ramosissima, 1647
salinity adaptation, 2083
Salmo marmoratus, 2711
Salmo trutta, 1003, 2523
salt tolerance, 1647
Salvelinus alpinus, 1017
Salvelinus umbla, 1017
Saxifraga oppositifolia, 1409
scale samples, 197
Scaptotrigona, 2107
scatology, 627
Scatophaga, 1781
Schistosoma mansoni, 1231
Scleractinia, 1339
scleractinian coral, 1177
scorpions, 925
sculpin, 1717
Scutellospora, 571
sea anemone, 2285
sea otter, 1899
Sebacinaceae, 1831
Sebastes inermis, 2591
secondary endosymbiotic bacteria, 2123
secondary sexual characters, 1533
selection, 113, 939, 1245, 1311, 1515, 2583
selfing, 2213
selfing rate, 879
sequence analysis, 2027
sex-assignment, 851
sex-biased dispersal, 125, 529
sex chromosomes, 2151
sex-markers, 851
sex ratio, 1275, 1533
sex ratio dynamics, 2151
sex-specific genetic structure, 1231
sexual reproduction, 1191
sexual selection, 1125
sexual size dimorphism, 1317
sibling species, 1427, 1453, 2065
Silurus glanis, 1039
simulated annealing, 2571
simulation, 1145, 2445
Sinapsis arvensis, 103
single nucleotide polymorphism, 603
SIT, 1915
size constraints, 2453
snakes, 1739
snowshoe hare, 1689
SNPs, 947
social behaviour, 507
social Hymenoptera, 2405
social insects, 1795
social organization, 1805
social structure, 289
sociality, 2307
soldier aphids, 1525
song, 473

Sorex araneus, 785
South America, 901
spatial autocorrelation, 305
spatial genetic structure, 725
spatial structure, 305
specialization, 191, 557
speciation, 619, 951, 1057, 1427, 1439, 1931, 2623
species boundaries, 1339
species concept, 1339
species discrimination, 437
sperm competition, 91
spotted eagle, 2189
SSCP, 1617, 1637, 1731, 2295
SSR function, 2453
SSRs, 1
SSU, 2669
SSU rRNA gene, 1555
stepping-stone colonization, 675
sterile insect technique, 1915
stingless bees, 2107
stocking, 1003
stocking impact, 1017
Stomatopod, 659
sturgeon, 1885
subdivided population, 2571
sublethal, 547
subspecies, 1739
substitution rate, 1905
subtropical tree, 2349
sunflowers, 1703
superinfections, 2425
SW Pacific, 1753
symbiosis, 17, 1565, 1845
sympatric speciation, 2337
Syncerus caffer, 267

Tapesia, 2229
Taraxacum, 253
tarsus length, 1317
temperate trees, 377
temporal change, 1515
temporal variation, 739
Tephritidae, 2425
Termitomyces, 1565
territory, 535
Tetraclita, 1731
Tetrao urogallus, 1669
Thamnophis sirtalis, 1739
Thaumetopoea pityocampa, 2435
thelytoky, 1815
Thymallus, 2599
Thymallus thymallus, 1393
Tomicus, 483
transgene cost, 1419
transgenic oilseed rape, 1419
trans-location, 421
transplant, 739
transposable elements, 347
trans-species evolution, 2157
Tricholoma, 557
Trigona, 2107

triple infection, 167
triploidy, 1499
*trn*L, 1957
trnL-F, 1965
Trollius europaeus, 2337
trophic, 2679
tropical forest, 2669
tropical rainforest, 725
tunicates, 1579
TWOGENER, 1657
Type III survivorship, 1029

Ursus americanus, 685
usurpation, 2405

variation, 17
Venezuela, 879
vertical transmission, 547
Vietnam, 1629
Vitis, 2013
Vouacapoua americana, 725

W chromosome, 851
walking-sticks, 1471
whales, 2241
whiteflies, 1219
wild rabbit, 2703
wild radish, 1419
wild turkey, 643
wild ungulates, 421
Wilsonia pusilla, 1605
Wolbachia, 167, 1275, 1815, 2425
worker policing, 1795
worker reproduction, 1795
wsp, 1815
wsp phylogeny, 167

Y-chromosome, 785
yolk protein electrophoresis, 2417

zooplankton, 1209
zygomycetes, 571